



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Invariant and quasi-invariant measures for Hamiltonian PDEs

Citation for published version:

Oh, T 2015, *Invariant and quasi-invariant measures for Hamiltonian PDEs*..

Link:

[Link to publication record in Edinburgh Research Explorer](#)

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



①

Invariant and quasi-invariant measures for Hamiltonian PDEs

Chap 1: Invariant Gibbs measures: Part 1.

Sec 1.1: Introduction

Finite dim'l Hamiltonian dynamics on \mathbb{R}^{2n}

$$(1) \left\{ \begin{array}{l} \partial_t p_{\hat{j}} = \frac{\partial H}{\partial q_{\hat{j}}} \\ \partial_t q_{\hat{j}} = - \frac{\partial H}{\partial p_{\hat{j}}} \end{array} \right\} = X \quad \hat{j} = 1, \dots, n.$$

with $H(p, q) = H(p_1, \dots, p_n, q_1, \dots, q_n)$

① Liouville's thm: given $\partial_t x = f(x)$,

$$\frac{d}{dt} \text{vol} = \text{div} f.$$

$$\begin{aligned} \Rightarrow \text{div} X &= \sum_{\hat{j}=1}^n \left[\frac{\partial}{\partial p_{\hat{j}}} X_{\hat{j}} + \frac{\partial}{\partial q_{\hat{j}}} X_{n+\hat{j}} \right] \\ &= \sum_{\hat{j}=1}^n \left[\frac{\partial}{\partial p_{\hat{j}}} \frac{\partial H}{\partial q_{\hat{j}}} + \frac{\partial}{\partial q_{\hat{j}}} \left(- \frac{\partial H}{\partial p_{\hat{j}}} \right) \right] = 0 \end{aligned}$$

\Rightarrow Lebesgue meas $dp dq = \prod_{\hat{j}=1}^n dp_{\hat{j}} dq_{\hat{j}}$ is invariant under (1).

(2)

② Hamiltonian H is conserved under (1).

$$\left(\begin{aligned} \frac{d}{dt} H(p(t), q(t)) \\ = \frac{\partial H}{\partial p} \frac{\partial p}{\partial t} + \frac{\partial H}{\partial q} \frac{\partial q}{\partial t} = 0. \end{aligned} \right.$$

\Rightarrow Gibbs measures:

$$d\mu = d\mu_\beta = Z^{-1} e^{-\beta H(p, q)} dp dq$$

are invariant under the dynamics (1). $\left(\begin{array}{l} \beta = \text{inv temp} \\ > 0 \end{array} \right.$

Invariance: $\Phi(t) : (p(0), q(0)) \mapsto (p(t), q(t))$.

$$\boxed{\mu(\Phi(t)A) = \mu(\{(p, q) \in \Phi(-t)A\})}$$

$$= \mu(\{\Phi(t)(p, q) \in A\})$$

$$= Z^{-1} \int_A \underbrace{e^{-\beta H(p(t), q(t))}}_{H(p(0), q(0))} \underbrace{dp(t) dq(t)}_{dp(0) dq(0)}$$

$$\boxed{= \mu(A)}$$

(3)

Rmk: Suppose that F is conserved under (1)

$$\Rightarrow d\mu_F = Z^{-1} e^{-F(p,q)} dp dq$$

is also invariant (for reasonable F).

Q: Why do we care about invariant meas?

① can view (1) as a dynamical system with meas-preserving transf $T := \Phi(1)$.

$$T : (p(0), q(0)) \mapsto (p(t), q(t)) \Big|_{t=1}$$

\Rightarrow Poincaré recurrence thm.

(i) \forall meas A with $\mu(A) > 0$,

$$\exists N \in \mathbb{N} \text{ s.t. } \mu(A \cap T^{-N}A) > 0.$$

(ii) a. e. $(p, q) \in \mathbb{R}^{2n}$ are stable according to Poisson. i.e.

$$\exists \{t_j\} \rightarrow \infty \text{ s.t.}$$

$$(p(t_j), q(t_j)) \rightarrow (p(0), q(0))$$

(4)

Also, Furstenberg's multiple recurrence thm

$\forall A$ with $\mu(A) > 0$ and $k \in \mathbb{N}$,

$\exists N \in \mathbb{N}$ s.t.

$$\mu(A \cap T^{-N}A \cap \dots \cap T^{-(k-1)N}A) > 0$$

(2) μ is "supposed" to describe a typical long time behavior of solutions.

Sec 1.2: Hamiltonian PDEs

- Nonlinear Schrödinger equation (NLS):

$$(NLS) \quad i\partial_t u + \Delta u = \pm |u|^{p-2} u \quad \text{on } \mathbb{T}^d.$$

- (generalized) Korteweg-de Vries equation (gKdV):

$$(gKdV) \quad \partial_t u + u_{xxx} = \pm u^{p-2} \partial_x u \quad \text{on } \mathbb{T}.$$

• $p=3$: KdV ($u \partial_x u$)

• $p=4$: modified KdV (mKdV with $u^2 \partial_x u$)

These are Hamiltonian PDEs with

$$H(u) = \frac{1}{2} \int_{\mathbb{T}^d} |\nabla u|^2 dx \pm \frac{1}{p} \int_{\mathbb{T}^d} |u|^p dx$$

(5)

$$(NLS) \iff \partial_t u = -i \frac{\partial H}{\partial \bar{u}}.$$

Symplectic space $L^2(\mathbb{T}^d)$

$$\text{symplectic form } \omega(f, g) = \text{Im} \int_{\mathbb{T}^d} f(x) \overline{g(x)} dx$$

$$\frac{dH}{du}(\phi) = \omega(\phi, -i \frac{\partial H}{\partial \bar{u}})$$

$$\uparrow \text{Gâteaux derivative} = \frac{d}{d\varepsilon} H(u + \varepsilon \phi) \Big|_{\varepsilon=0}.$$

On the Fourier side:

$$p_n = \text{Re } \hat{u}_n, \quad q_n = \text{Im } \hat{u}_n.$$

$$H(p, q) = \frac{1}{2} \sum |m|^2 (p_n^2 + q_n^2)$$

$$(NLS) \quad \partial_t u = i \Delta u + \text{nonlin}$$

$$\iff \partial_t \hat{u}_n = -i |m|^2 \hat{u}_n$$

$$\iff \partial_t \begin{pmatrix} p_n \\ q_n \end{pmatrix} = \begin{pmatrix} \frac{\partial H}{\partial q_n} \\ -\frac{\partial H}{\partial p_n} \end{pmatrix}.$$

(6)

Note that $H(u)$ is conserved under the dynamics.

Moreover, we use the conservation of mass

$$M(u) = \int |u|^2 dx.$$

Natural question:

Is the Gibbs measure

$$d\mu = Z^{-1} e^{-\beta H(u)} du$$

invariant under NLS (or gKdV)?

$\beta=1$

$$\begin{aligned} d\mu &= Z^{-1} e^{-H(u)} du \\ &= Z^{-1} e^{-\frac{1}{p} \int |u|^p dx} \underbrace{e^{-\frac{1}{2} \int |\nabla u|^2 dx}}_{\text{Wiener measure on } \Pi^d} du \end{aligned}$$

Wiener measure on Π^d .

In the following, we consider

$$d\mu = Z^{-1} e^{-H(u) - \frac{1}{2} M(u)} du$$

for NLS in order to avoid the problem at freq 0.

(7)

Sec 1.3: Gaussian measures on H^s .

Consider

$$d\rho_s = Z^{-1} e^{-\frac{1}{2} \|u\|_{H^s}^2} du$$

as a limit of

$$d\rho_{s,N} = Z_N^{-1} e^{-\frac{1}{2} \|P_{\leq N} u\|_{H^s}^2} dP_{\leq N} u$$

$$= Z_N^{-1} \prod_{|n| \leq N} \underbrace{e^{-\frac{1}{2} \langle n \rangle^{2s} |\hat{u}_n|^2}}_{\text{Gaussian meas on } \mathbb{C}} d\hat{u}_n$$

Gaussian meas on \mathbb{C} .
mean 0, var = 2 $\langle n \rangle^{-2s}$.

Namely, $\rho_{s,N}$ is the induced prob meas under the map

$$\omega \longmapsto u^N = \sum_{|n| \leq N} \frac{g_n(\omega)}{\langle n \rangle^s} e^{in \cdot x}$$

where $\{g_n\}$ = indep. standard \mathbb{C} -valued Gaussian r.v.'s.

$$\mathbb{E}(g_n) = 0, \text{ Var}(g_n) = 2.$$

Problem: We can not take a limit $N \rightarrow \infty$

in $H^s(\mathbb{T}^d)$!!

⑧

Let $\sigma \in \mathbb{R}$ and $M > N$.

$$\mathbb{E} \left[\|u^M - u^N\|_{H^\sigma(\mathbb{T}^d)}^2 \right]$$

$$= \mathbb{E} \sum_{N < |n| \leq M} \frac{|g_n|^2}{\langle n \rangle^{2s-2\sigma}}$$

$$\sim \sum_{N < |n| \leq M} \frac{1}{\langle n \rangle^{2s-2\sigma}} \rightarrow 0$$

if and only if $2s - 2\sigma > d$

$$\Leftrightarrow \sigma < s - \frac{d}{2}$$

(2)

If (2) is satisfied, then

u^N converges in $L^2(\Omega; H^\sigma(\mathbb{T}^d))$.

and define ρ_s as the induced prob meas under

$$\omega \mapsto u = \sum_{n \in \mathbb{Z}^d} \frac{g_n}{\langle n \rangle^s} e^{in \cdot x}$$

\Leftarrow prob meas on $H^\sigma(\mathbb{T}^d)$

Lec Notes in Math.
Gross '65, Kuo '75

Abstract Wiener space: $H = \infty$ -dim'l Hilbert space ^{separable} ⑨
" $df = z^{-1} e^{-\frac{1}{2} \|u\|_H^2} du$ "

\Leftarrow NOT countably additive if $\dim H = \infty$.

\Rightarrow Must work on a larger space $B \supset H$ ^{densely & conti.}
 \uparrow
separable Banach space

We say (B, H, μ) is an abstract Wiener space

if $\int_B e^{i\langle u, \varphi \rangle} d\mu(u) = e^{-\frac{1}{2} \|\varphi\|_{H^*}^2}$
 \uparrow
 $B-B^*$ duality.

for all $\varphi \in B^* \subset H^*$.

Rmk: ① $i: H \hookrightarrow B$.

The triplet (i, H, B) is often called an abstract Wiener space.

② $B =$ completion of H under a "measurable" seminorm $\|\cdot\|_B$:

$\forall \varepsilon > 0 \exists P_\varepsilon \in \mathcal{F} =$ collection of
finite dim'l ortho proj.

s.t.

$$P(\|P u\|_B > \varepsilon) < \varepsilon$$

for all $P \in \mathcal{F}$ orthogonal to P_ε .

ex: $H = H^s$

$$B = H^\sigma, \quad \sigma < s - \frac{d}{2}$$

$$W^{\sigma,p}$$

$$\mathcal{F}L^{\sigma,p}, \quad (\sigma - s)p < -d$$

$$\|f\|_{\mathcal{F}L^{\sigma,p}(\mathbb{T}^d)} = \|\langle n \rangle^\sigma \hat{f}_n\|_{\ell^p(\mathbb{Z}^d)}$$

Lemma 1.1 (Tail estimate) Let $\sigma < s - \frac{d}{2}$.

Then, $\rho_\sigma(\{\|u\|_{H^\sigma} > K\}) \leq c e^{-cK^2}$

for all $K > 0$.

Rmk: This follows from Fernique's integrability thm:

$$\int_B e^{c\|u\|_B^2} d\rho(u) < \infty.$$

Pf: By Chebyshev's ineq,

$$\begin{aligned} e^{cK^2} \rho_\sigma(B_K^c) &\leq \int_{H^\sigma} e^{c\|u\|_{H^\sigma}^2} d\rho_\sigma(u) \\ &= \prod_{n \in \mathbb{Z}^d} \int_{\mathbb{C}} e^{c\langle n \rangle^{2\sigma-2s} |g_n|^2} e^{-\frac{1}{2}|g_n|^2} \frac{dg_n}{2\pi} \end{aligned}$$

(11)

$$= \prod_{n \in \mathbb{Z}^d} \frac{1}{1 - 2C \langle n \rangle^{2\sigma - 2s}}$$

$$\left(\begin{array}{c} \mathbb{E}[e^{aX^2}] = (1 - 2a)^{-1/2}, \quad a < 1/2 \\ \uparrow \\ X \sim N_{\mathbb{R}}(0, 1) \end{array} \right.$$

$$= \prod_{n \in \mathbb{Z}^d} \left(1 + \frac{2C \langle n \rangle^{2\sigma - 2s}}{1 - 2C \langle n \rangle^{2\sigma - 2s}} \right)$$

$$< \infty \quad \text{iff} \quad \sigma < s - \frac{d}{2}$$

$$\left(\begin{array}{l} a_n > 0. \text{ Then} \\ \prod (1 + a_n) < \infty \Leftrightarrow \sum a_n < \infty \end{array} \right.$$



Sec 1.4: Construction of Gibbs measure on \mathbb{T}

Fix $d=1$:

$$\begin{aligned} d\mu &= Z^{-1} e^{-H(u) - \frac{1}{2} M(u)} du \\ &= Z^{-1} e^{-\frac{1}{p} \int |u|^p dx} d\rho \end{aligned}$$

where

$\rho = \rho_1 =$ Wiener meas on \mathbb{T} .

Under ρ , we have \uparrow actually, o-v.

(3) $u = \sum_n \frac{g_n}{\langle n \rangle} e^{inx}.$

Compare this with Fourier-Wiener series of a Brownian motion on $[0, 2\pi)$:

$$B(t) = g_0 t + \sum_{n \neq 0} \frac{g_n}{\sqrt{2\pi} i n} e^{int} - \sum_{n \neq 0} \frac{g_n}{\sqrt{2\pi} i n}$$

Defocusing case (with - sign):

$$(3) \Rightarrow u \in H^s(\mathbb{T}) \quad s < \frac{1}{2} \text{ a.s.}$$

$$\Rightarrow u \in L^p(\mathbb{T}), \quad p < \infty \text{ a.s.}$$

Sobolev

$$\Rightarrow 0 < e^{-\frac{1}{p} \int |u|^p dx} \leq 1 \quad \text{a.s.}$$

$\Rightarrow \mu$ is a prob meas on $H^0(\pi) \setminus H^{1/2}(\pi)$
 $\mu \ll \rho.$ $s < 1/2.$

• Focusing case: Let $p > 2.$

$$\begin{aligned} \int_{\pi} |u|^p &\geq \|u\|_{L^2}^p = \left(\sum |\hat{u}_n|^2 \right)^{p/2} \\ &\geq \sum \left| \frac{g_n}{\langle n \rangle} \right|^p \quad (l^2 \subset l^p) \end{aligned}$$

$$\Rightarrow \mathbb{E}_A \left[e^{\int |u|^p} \right] \geq \prod_n \mathbb{E} \left[e^{|g_n|^p / \langle n \rangle^p} \right] = \infty.$$

$$\left(\begin{aligned} \mathbb{E}[e^{|g|^p}] &= \frac{1}{2\pi} \int e^{|g|^p} e^{-|g|^2/2} dg \\ &= \infty. \end{aligned} \right.$$

• Remedy: Mass is conserved.

\Rightarrow Introduce a mass cutoff.

Prop 1.2: (Lebowitz-Rose-Speer '88, Bourgain '94)

(ii) $p < 6$

(4) $R(u) = R_r(u) := \mathbb{1}_{\{\|u\|_{L^2} \leq r\}} e^{\frac{1}{r} \int \|u\|^p} \in L^q(d\rho)$

$$\forall r, q < \infty.$$

(ii) $p = 6$: (4) holds if r is sufficiently small.

(In fact, if $r < \|Q\|_{L^2(\mathbb{R})}$
 ↑ ground state
 related to (minimal mass) finite time blowup solution

(iii) If $p > 6$ or $p = 6$ and $r \geq \|Q\|_{L^2(\mathbb{R})}$, then (4) fails.

Rmk: $p=6$: mass-critical NLS

\Leftarrow lowest power for which we have finite time blowups.

Lemma 1.3: $\{\tilde{g}_n\}$ indep standard \mathbb{R} -valued Gaussian r.v.'s.

Then,

$$P\left[\left(\sum_{n=1}^M \tilde{g}_n^2\right)^{1/2} \geq R\right] \leq e^{-\frac{1}{4}R^2}, \quad R \geq 3M^{1/2}, \quad M \geq 1$$

Pf: $0 < t < 1/2$

$$\begin{aligned} (\text{LHS}) &\leq e^{-tR^2} \mathbb{E}\left[e^{t \sum_{n=1}^M \tilde{g}_n^2}\right] \\ &= (1-2t)^{-M/2} e^{-tR^2}. \end{aligned}$$

Choose $t = \frac{1}{2} \left(1 - \frac{M}{R^2}\right)$.

$$\Rightarrow (\text{LHS}) = \left(\frac{R^2}{M}\right)^{M/2} e^{-\frac{1}{2}R^2 + \frac{1}{2}M}$$

$$\leq e^{\frac{M}{2} \log \frac{R^2}{M} + \left(\frac{1}{18} - \frac{1}{2}\right)R^2} \leq e^{-\frac{1}{4}R^2}$$

$$\left(\log x \leq \frac{x}{4} \text{ for } x \geq 9 \right)$$

□

Pf of Prop 1.2 (i) and (ii):

(16)

• By Bernstein's ineq,

$$\|P_{\leq M_0} u\|_{L^p} \leq c M_0^{\frac{1}{2}-\frac{1}{p}} \underbrace{\|P_{\leq M_0} u\|_{L^2}}_{\leq r}$$

$\lambda \geq 1$ Choose M_0 s.t.

$$\frac{1}{2} \lambda = c M_0^{\frac{1}{2}-\frac{1}{p}} r$$

i.e.

(5) $M_0 \sim \left(\frac{\lambda}{r}\right)^{\frac{1}{\frac{1}{2}-\frac{1}{p}}}$

• Let $M_j = 2^j M_0$. Then, we have

$$\|P_{M_j} u\|_{L^p} \leq c M_j^{\frac{1}{2}-\frac{1}{p}} \|P_{M_j} u\|_{L^2}.$$

• Let $\{\sigma_j\}$ s.t. $\sum_{j \geq 1} \sigma_j = \frac{1}{2}$.

$$\Leftarrow \text{set } \sigma_j = C 2^{-\varepsilon j} = C M_0^\varepsilon M_j^{-\varepsilon}.$$

By dyadic pigeon hole principle,

$$\begin{aligned} & \mathcal{P}(\|u\|_{L^p} > \lambda, \|u\|_{L^2} \leq r) \\ & \leq \sum_{j=1}^{\infty} \underbrace{\mathcal{P}(\|P_{M_j} u\|_{L^p} > \sigma_j \lambda)}_{\leq \mathcal{P}(\|P_{M_j} u\|_{L^2} > c \sigma_j M_j^{\frac{1}{2}-\frac{1}{p}} \lambda)} \end{aligned}$$

(17)

$$P \left(\|P_{M_j} u\|_{L^2} > c \sigma_j M_j^{\frac{1}{p}-\frac{1}{2}} \lambda \right) \\ \sim P \left(\left(\sum_{|n| \sim M_j} |g_n(\omega)|^2 \right)^{\frac{1}{2}} \gtrsim \underbrace{\sigma_j M_j^{\frac{1}{p}+\frac{1}{2}} \lambda}_{R_j} \right)$$

$$\begin{aligned} R_j &:= \sigma_j M_j^{\frac{1}{p}+\frac{1}{2}} \lambda & \lambda \geq 1 \\ &\gtrsim M_0^\varepsilon M_j^{\frac{1}{2}+\frac{1}{p}-\varepsilon} \\ &\gtrsim M_0^\varepsilon M_j^{\frac{1}{2}+}. \end{aligned}$$

Lemma 1.3.

$$\Rightarrow P(\|u\|_{L^p} > \lambda, \|u\|_{L^2} \leq r)$$

$$\begin{aligned} &\leq \sum_{j=1}^{\infty} e^{-c R_j^2} \\ &\sim e^{-c \lambda^2 M_0^{\frac{2}{p}+1}} \quad R_j^2 \sim M_0^{\frac{2}{p}+1} \lambda^2 2^{(\frac{2}{p}+1-2\varepsilon)j} \end{aligned}$$

$$\begin{aligned} (5) \quad &\sim e^{-c \lambda^{\frac{4p}{p-2}} r^{\frac{-2p+4}{p-2}}} \lesssim \begin{cases} e^{-c \lambda^{p+s}}, & p > 6 \\ e^{-c \lambda^p} & \text{with } c \gg 1 \text{ if } p = 6 \end{cases} \\ &\frac{4p}{p-2} > p \text{ for } p < 6 \end{aligned}$$

□

$$R_r(u) = \frac{1}{\{\|u\|_{L^2} \leq r\}} e^{\frac{1}{p} \int |u|^p}$$

$$R_{N,r}(u) = \frac{1}{\{\|P_N u\|_{L^2} \leq r\}} e^{\frac{1}{p} \int |P_N u|^p}$$

Cor 1.4: $\forall q < \infty$,
 $R_{N,r}(u) \rightarrow R_r(u)$ in $L^q(df)$
 as $N \rightarrow \infty$.

In particular,

$$d\mu_{N,r} = Z_N^{-1} R_{N,r}(u) df$$

converges "uniformly" to $\mu = \mu_r$.

i.e. $\forall \varepsilon > 0$, $\exists N_0 \in \mathbb{N}$ s.t.

$$|\mu_{N,r}(A) - \mu(A)| < \varepsilon$$

$\forall N \geq N_0$ and measurable A .

pf: $\int |P_{\Sigma N} u|^p \rightarrow \int |u|^p \quad \text{a.s.}$

$\Rightarrow R_{N,r}(u) \rightarrow R_r(u) \quad \text{a.s.}$

\Rightarrow By Egoroff's thm,

$R_{N,r}(u) \rightarrow R_r(u)$ almost uniformly
 \Rightarrow in measure.

• Given $\varepsilon > 0$,

let $A_{N,\varepsilon} = \left\{ u \in H^{\frac{1}{2}}(\Pi) : |R_{N,r}(u) - R_r(u)| \leq \frac{1}{2}\varepsilon \right\}$.

Then, $\rho(A_{N,\varepsilon}^c) \rightarrow 0$.

$\Rightarrow \|R_{N,r} - R_r\|_{L^q(\rho)}$

$\leq \| (R_{N,r} - R_r) \mathbb{1}_{A_{N,\varepsilon}} \|_{L^q}$

$+ \| (R_{N,r} - R_r) \mathbb{1}_{A_{N,\varepsilon}^c} \|_{L^q}$

$\leq \frac{1}{2}\varepsilon + \underbrace{\left(\|R_{N,r}\|_{L^{2q}} + \|R_r\|_{L^{2q}} \right)}_{\leq C < \infty} \left\{ \rho(A_{N,\varepsilon}^c) \right\}^{\frac{1}{2p}}$

$< \varepsilon$.



Sec 1.5: Invariance of the Gibbs measure.

(20)

3 scenarios.

• Case I: We have an a priori (deterministic) global well-posedness in $\text{supp}(\mu) \supset H^{\frac{1}{2}-}(\mathbb{T})$.

ex: KdV, cubic NLS.

\Rightarrow easy

Case II: Only a priori local-in-time dynamics.

Idea: use invariance (in place of a conservation law) to construct global-in-time dynamics



Bourgain '94: use invariance of the finite dim'l

Gibbs meas associated to the finite dim'l approx

\Rightarrow a.s. GWP. \Rightarrow invariance.

Case III: No local well-posedness.

(III.1) probabilistic local-in-time Cauchy theory
Bourgain '96, Burq-Tzvetkov '07

(III.2) compactness argument
(for measures on space-time functions)

Burq-Thomann-Tzvetkov '14, Oh-Thomann '15.

In the following, we focus on Case II.

$$(NLS) \quad i\partial_t u + \Delta u = \pm |u|^{p-2} u.$$

Suppose that we have subcritical LWP in $\text{supp}(u)$.

(\Leftarrow Bourgain '93)

i.e. time of local existence $\delta \sim (1 + \|u_0\|_B)^{-\theta}$

$$B = H^{\frac{1}{2}-\varepsilon}(\mathbb{T})$$

Then, the same argument yields LWP of the following "finite dimensional" approx

$$(FNLS) \quad \begin{cases} i\partial_t u^N + \Delta u^N = \pm P_{\leq N}(|u^N|^{p-2} u^N) \\ u^N|_{t=0} = u_0 \end{cases}$$

for $|t| \leq \delta$.

• (FNLS) is NOT finite dim'l.

$$\cdot \partial_t u^N = -i \frac{\partial H_N}{\partial \overline{u^N}}$$

$$\text{where } H_N(u^N) = \frac{1}{2} \int |\partial_x u^N|^2 \pm \frac{1}{p} \int |P_{\leq N} u^N|^p$$

↑

lin. dynamics on high freq.

① $v^N := P_{\leq N} u^N$ satisfies

$$(FNLS_{\text{low}}) \quad i \partial_t v^N + \partial_x^2 v^N = \pm P_{\leq N} (|v^N|^{p-2} v^N)$$

finite dim'l system of ODEs (on the Fourier side)

② $P_{>N} u^N$ evolves linearly:

$$(FNLS_{\text{high}}) \quad \partial_t \widehat{u^N}(n) = -i n^2 \widehat{u^N}(n), \quad |n| > N.$$

• As a lin. eqn, $(FNLS_{\text{high}})$ is GWP.

• $(FNLS_{\text{low}})$ preserves the L^2 -norm:

$$\|v^N\|_{L^2} = \left(\sum_{|n| \leq N} |\widehat{v^N}(n)|^2 \right)^{1/2}$$

= Euclidean distance on \mathbb{C}^{2N+1}

$\Rightarrow (FNLS_{\text{low}})$ is GWP.

⇒ (FNLS) is GWP for each $N \in \mathbb{N}$.

(23)

but there is NO uniform (in N) control on $\|u^N(t)\|_B$.

Write $\mathcal{F} = \mathcal{F}_N \otimes \mathcal{F}_N^\perp$

$$\begin{cases} d\mathcal{F}_N = Z_N^{-1} e^{-\frac{1}{2} \|P_{\leq N} u\|_{H'}^2} dP_{\leq N} u \\ d\mathcal{F}_N^\perp = Z_N^{-1} e^{-\frac{1}{2} \|P_{> N} u\|_{H'}^2} dP_{> N} u \end{cases}$$

Then,

① $d\mu_{N, \text{low}} = Z_N^{-1} R_{N, r}(v^N) d\mathcal{F}_N$ is invariant under (FNLS_{low}).

② \mathcal{F}_N^\perp is invariant under (FNLS_{high}).

⇒ $\mu_N := \mu_{N, \text{low}} \otimes \mathcal{F}_N^\perp$ is invariant under (FNLS).

Prop 1.5 (Bourgain '94)

(24)

$\forall T < \infty, \varepsilon > 0, \exists \Omega_N = \Omega_N(T, \varepsilon)$ s.t.

(i) $\mu_N(\Omega_N^c) < \varepsilon$

(ii) For $u_0 \in \Omega_N$, $\exists!$ soln u^N to (FNLS)

s.t.

$s \leq \frac{1}{2}$ $\|u^N(t)\|_{H^s} \lesssim \left(\log \frac{T}{\varepsilon}\right)^{\frac{1}{2}}, \quad |t| \leq T.$

Pf: Local theory:

$$\|u_0\|_{H^s} \leq K \Rightarrow \|u^N(t)\|_{H^s} \leq 2K$$

$$\text{for } |t| \leq \delta \sim K^{-\theta}.$$

Let $\Omega_N = \bigcap_{j=-[T/\delta]}^{[T/\delta]} \Phi_N(j\delta) \left(\|u_0\|_{H^s} \leq K \right)$

(i) By invariance of μ_N ,

$$\mu_N(\Omega_N^c) \lesssim \frac{T}{\delta} \mu_N(\Phi_N(j\delta)(B_K^c))$$

$$\lesssim T K^\theta e^{-cK^2} < \varepsilon$$



Prop 1.2 & Lemma 1.1.

by choosing $K \sim \left(\log \frac{T}{\varepsilon}\right)^{\frac{1}{2}}.$

(ii) By construction,

$$\|u^N(j\delta)\|_{H^s} \leq K, \quad j = 0 \neq 1, \dots, \pm [T/\delta]$$

\Rightarrow By local theory,

$$\|u^N(t)\|_{H^s} \leq 2K \sim \left(\log \frac{T}{\varepsilon}\right)^{1/2}, \quad |t| \leq T$$

□

Lemma 1.6 (Approximation Lemma) $s < 1/2$

$u_0 \in H^s$ with $\|u_0\|_{H^s} \leq K$.

Suppose soln u^N to (FNLS) with $u^N|_{t=0} = u_0$ satisfies

$$\|u^N(t)\|_{H^s} \leq K, \quad |t| \leq T.$$

Then, $\exists!$ soln u to (NLS) on $[-T, T]$ with $u|_{t=0} = u_0$.

Moreover, $\exists C_0, C_1, C_2$ s.t.

$$\|u(t) - P_N u^N(t)\|_{H^{s_1}} \leq C_0 e^{C_1(1+K)^{C_2} T} K \underbrace{N^{s_1-s}}_{\rightarrow 0}$$

for $s_1 < s$.

Pf: (FNLS) and (NLS) with $u^N|_{t=0} = u|_{t=0} = u_0$ (26)
 are locally well-posed on $[-\delta, \delta]$, $\delta \sim (1+K)^{-\theta}$
 indep of N .

\Rightarrow With $v^N = P_{\leq N} u^N$,

$$\|u - v^N\|_{X^{s_1, b}}$$

$$\lesssim \underbrace{\| \eta(t) \delta(t) P_{>N} u \|_{H^{s_1}}}_{\substack{\uparrow \\ \text{smooth cutoff}}} + \text{nonlin. terms}$$

$$\lesssim K N^{s_1 - s}$$

$$\textcircled{1} (Id - P_{\leq N}) (|u|^{p-2} u)$$

$$\lesssim \delta^\theta N^{s_1 - s} K^{p-1}$$

$$+ \textcircled{2} P_{\leq N} (|u|^{p-2} u - |v^N|^{p-2} v^N)$$

$$\lesssim \delta^\theta K^{p-2} \|u - v^N\|_{X^{s_1, b}}$$

$$\Rightarrow \|u - v^N\|_{X^{s_1, b, \delta}} \leq C K N^{s_1 - s} + \frac{1}{2} \|u - v^N\|_{X^{s_1, b, \delta}}$$

$$\Rightarrow \|u - v^N\|_{X^{s_1, b, \delta}} \leq 2C K N^{s_1 - s}$$

Repeating $\sim T/\delta$ many times, we obtain

$$\|u(t) - v^N(t)\|_{H^{s_1}} \lesssim \underbrace{e^{T/\delta}}_{\rightarrow e^{C_1(1+K)T}} K N^{s_1 - s} e^{C_2 T} \quad \square$$

Prop 1.7 (Almost a.s. GWP) $s < 1/2$ (27)

Given $T, \varepsilon > 0$, $\exists \Omega_{T, \varepsilon}$ s.t.

(i) $\mu(\Omega_{T, \varepsilon}^c) < \varepsilon$

(ii) For $u_0 \in \Omega_{T, \varepsilon}$, $\exists!$ soln u to (NLS) on $[-T, T]$

s.t. $\|u(t)\|_{H^s} \lesssim \left(\log \frac{1}{\varepsilon}\right)^{1/2}, |t| \leq T.$

Pf:

Let $\Omega_N(T, \varepsilon)$ be as in Prop 1.5.

From Prop 1.5, $\|\Phi_N(t)(u_0)\|_{H^s} \leq 2K$

for $|t| \leq T$ and $u_0 \in \Omega_N$.

• By Lemma 1.6, $\exists N_1 \in \mathbb{N}$ s.t.

$$\|u(t) - u^N(t)\|_{H^{s_1}} \ll 1, \quad |t| \leq T$$

for $N \geq N_1$

$$\Rightarrow \|u(t)\|_{H^{s_1}} \lesssim K \sim \left(\log \frac{1}{\varepsilon}\right)^{1/2}, \quad |t| \leq T$$

• Also, $\mu(\Omega_N^c) \stackrel{\text{c.s.}}{\lesssim} \left(\int_{\Omega_N^c} \underbrace{\mathbb{1}_{\{\|u\|_{L^2} \leq r\}}}_{\leq \mathbb{1}_{\{\|P_{\leq N} u\|_{L^2} \leq r\}}} d\mu \right)^{1/2}$

$$\stackrel{\text{c.s.}}{\lesssim} \left(\mu_N(\Omega_N^c) \right)^{1/4} < \varepsilon.$$

□

Thm 1.8: NLS is a.s. GWP with respect to μ .

(28)

Pf: Given $\varepsilon > 0$, let $T_j = 2^j$, $\varepsilon_j = \frac{\varepsilon}{2^j}$

\Rightarrow Construct $\Omega_j = \Omega_{T_j, \varepsilon_j}$.

• Let $\Omega_\varepsilon = \bigcap_{j=1}^{\infty} \Omega_j$

$$(i) \mu(\Omega_\varepsilon^c) < \sum_{j=1}^{\infty} \frac{\varepsilon}{2^j} = \varepsilon$$

(ii) If $u_0 \in \Omega_\varepsilon$, then $\exists!$ soln u on $[-T_j, T_j]$
for any $j \in \mathbb{N} \Rightarrow$ global soln.

• Let $\Sigma = \bigcup_{\varepsilon > 0} \Omega_\varepsilon$

$$(i) \mu(\Sigma^c) = \inf_{\varepsilon > 0} \varepsilon = 0.$$

(ii) If $u_0 \in \Sigma$, then $u_0 \in \Omega_\varepsilon$ for some $\varepsilon > 0$.
 $\Rightarrow \exists!$ global soln u with $u|_{t=0} = u_0$

□

Thm 1.9: The Gibbs measure μ is invariant under NLS.

Pf: By time reversibility of $\Phi(t)$, it suffices to show

(b) $\mu(A) \leq \mu(\Phi(t)A)$

for all measurable set $A \subset \Sigma$
and $t \in \mathbb{R}$

By inner regularity, we have

$$\mu(A) = \sup_{\substack{F \subset A \\ \text{closed in } H^s, s < 1/2}} \mu(F)$$

i.e. $\exists \{F_n\}$ closed in H^s s.t. $F_n \subset A$ and

$$\mu(A) = \lim_{n \rightarrow \infty} \mu(F_n)$$

- Suffices to prove (b) for closed sets.

$$\begin{aligned} \text{Then, } \mu(A) &= \lim_{n \rightarrow \infty} \mu(F_n) \\ &\leq \overline{\lim}_{n \rightarrow \infty} \mu(\Phi(t)F_n) \\ &\leq \mu(\Phi(t)A) \\ &\quad \uparrow F_n \subset A. \end{aligned}$$

- Given a closed set $F \subset H^s$,

$$\text{let } K_n = \{u \in F : \|u\|_{H^\sigma} \leq n\}, \quad s < \sigma < 1/2.$$

Then, K_n is compact in H^s .

\Rightarrow Suffices to prove (b) for compact sets.

$$\text{Then, } \mu(F) = \lim_{n \rightarrow \infty} \mu(K_n) \quad \text{Lemma 1.1 \& Prop 1.2}$$

$$\leq \overline{\lim}_{n \rightarrow \infty} \mu(\Phi(t)K_n)$$

$$\leq \mu(\Phi(t)F)$$

$$K_n \subset F$$

Assume
hold. in H^0

30

Let K be a compact set in H^1 .

Since $\mu_N \rightarrow \mu$ & Portmanteau thm, we have

$$(7) \quad \mu(\Phi(t)K + \overline{B_\varepsilon}) \geq \overline{\lim} \mu_N(\Phi(t)K + \overline{B_\varepsilon})$$

Fix $t \ll 1$. Then,

$$\Phi_N(t)(K + B_\delta) \subset \Phi_N(t)K + B_{\varepsilon/2}$$

$$\subset \Phi(t)K + B_\varepsilon$$

↑ Approximation lemma (Lemma 1.6)

⇒ By invariance of μ_N ,

$$(8) \quad \mu_N(K + B_\delta) = \mu_N(\Phi_N(t)(K + B_\delta)) \\ \leq \mu_N(\Phi(t)K + B_\varepsilon)$$

Hence,

$$\mu(K) \leq \mu(K + B_\delta)$$

$$\leq \underline{\lim} \mu_N(K + B_\delta)$$

$$\stackrel{(8)}{\leq} \underline{\lim} \mu_N(\Phi(t)K + B_\varepsilon)$$

$$\leq \overline{\lim} \mu_N(\Phi(t)K + \overline{B_\varepsilon})$$

$$\stackrel{(7)}{\leq} \mu(\Phi(t)K + \overline{B_\varepsilon})$$

$$\Rightarrow \text{Let } \varepsilon \rightarrow 0. \quad \mu(K) \leq \mu(\Phi(t)K).$$

□

Chap 2: Invariant Gibbs measures, Part 2. (31)

Sec 2.1: Gibbs measures on \mathbb{T}^2 .

We would like to construct

$$\begin{aligned} d\mu &= Z^{-1} e^{-H(u)} du \\ &= Z^{-1} e^{-\frac{1}{p} \int |u|^p} d\rho \quad \text{on } \mathbb{T}^2. \end{aligned}$$

- ρ is supported on $H^s(\mathbb{T}^2) \setminus L^2(\mathbb{T}^2)$, $s < 0$.

$$\Rightarrow \int_{\mathbb{T}^2} |u|^p dx = \infty \quad \text{a.s.}$$

\Rightarrow We need to renormalize $\int |u|^p$

- In the following, we focus on the defocusing case.

(

- Focusing case: Brydges - Slade '96.
We cannot construct Gibbs meas for the focusing cubic NLS even if we consider the Wick ordered nonlinearity (with the Wick ordered L^2 -cutoff.)

)

Sec 2.2: Hermite polynomials, white noise functional, and Wick ordering

(32)

Hermite poly $H_k(x; \sigma)$ defined through the generating function

$$G(t, x; \sigma) = e^{tx - \frac{1}{2}\sigma t^2} \\ = \sum_{k=0}^{\infty} \frac{t^k}{k!} H_k(x; \sigma)$$

$$H_0(x; \sigma) = 1, \quad H_1(x; \sigma) = x \\ H_2(x; \sigma) = x^2 - \sigma, \quad H_3(x; \sigma) = x^3 - 3\sigma x, \\ H_4(x; \sigma) = x^4 - 6\sigma x^2 + 3\sigma^2$$

• White noise functional

$$W(\cdot): L^2(\mathbb{T}^2) \rightarrow L^2(\mathcal{Q})$$

$$\text{by } W_f(\omega) = \langle f, w(\omega) \rangle_{L^2} = \sum_{n \in \mathbb{Z}^2} \hat{f}(n) \overline{g_n(\omega)}$$

$$\text{where } w = \sum g_n(\omega) e^{in \cdot x} \leftarrow \text{white noise on } \mathbb{T}^2$$

$\Rightarrow W_f$ is a Gaussian r.v with mean 0
var $\|f\|_{L^2}^2$

Moreover,

$$E[W_f \overline{W_h}] = \langle f, h \rangle_{L^2}$$

For simplicity, we work in the real-valued ⁽³³⁾ setting. i.e. $g_{-n} = \overline{g_n}$, etc.

$$\text{Var}(g_n) = 1$$

See Oh-Thomann '15 for \mathbb{C} -valued setting involving the (generalized) Laguerre polynomials.

Lemma 2.1 $f, h \in L^2(\mathbb{T}^2)$ with $\|f\|_{L^2} = \|h\|_{L^2} = 1$.

Then, we have

$$\mathbb{E} [H_k(W_f) H_m(W_h)] = \delta_{km} k! [\langle f, h \rangle_{L^2}]^k$$

Pf.: First, recall

$$\int_{\mathbb{R}} e^{W_f(\omega)} d\rho = \prod_{n \in \mathbb{I}^*} \frac{1}{\pi} \int_{\mathbb{C}} e^{2\text{Re}(\hat{f}_n \overline{g_n}) - |g_n|^2} dg_n$$

$$\times \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{\hat{f}_0 g_0} e^{-\frac{1}{2} g_0^2} dg_0$$

$$= \prod_{n \in \mathbb{I}^*} \frac{1}{\pi} \int_{\mathbb{R}} e^{2\text{Re} \hat{f}_n \text{Re} g_n - (\text{Re} g_n)^2} d\text{Re} g_n \int_{\mathbb{R}} e^{2\text{Im} \hat{f}_n \text{Im} g_n - (\text{Im} g_n)^2} d\text{Im} g_n$$

$$\times \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \dots dg_0$$

$$= \sum_{n \in \mathbb{I}^*} |\hat{f}_n|^2 e^{\frac{1}{2} (\hat{f}_0)^2} = e^{\frac{1}{2} \|f\|_{L^2}^2}.$$

$$\mathbb{I} = \text{index set} = " \mathbb{Z}^2 / 2 " = \mathbb{Z} \times \mathbb{N} \cup \mathbb{N} \times \{0\} \cup \{(0,0)\}$$

$$\mathbb{I}^* = \mathbb{I} \setminus \{(0,0)\}$$

$$\begin{aligned}
&\Rightarrow \int_{\Omega} G(t, W_f(\omega)) G(s, W_h(\omega)) dP(\omega) \\
&= e^{-\frac{1}{2}(t^2 + s^2)} \int_{\Omega} e^{W_{tf+sh}(\omega)} dP(\omega) \\
&= e^{ts \langle f, h \rangle_{L^2}} e^{\frac{1}{2} \|tf + sh\|_{L^2}^2}
\end{aligned}$$

$$\Rightarrow e^{ts \langle f, h \rangle_{L^2}} = \sum_{k, m=0}^{\infty} \frac{t^k s^m}{k! m!} \int_{\Omega} H_k(W_f) H_m(W_h) dP$$

$$\sum_{k=0}^{\infty} \frac{(ts)^k}{k!} (\langle f, h \rangle)^k.$$

□

Under f , we have

$$U = \sum_{n \in \mathbb{Z}^2} \frac{g_n}{\langle n \rangle} e^{in \cdot x} \quad \begin{array}{l} g_{-n} = \overline{g_n} \\ \text{Var}(g_n) = 1 \end{array}$$

Given $N \in \mathbb{N}$, let $U_N := P_{|n| \leq N} U$

Then, $U_N(x)$ is a mean 0 \mathbb{R} -valued Gaussian r.v.'s with variance

$$\sigma_N := \mathbb{E}[U_N^2(x)] = \sum_{|n| \leq N} \frac{1}{\langle n \rangle^2} \sim \log N.$$

Define the Wick ordered monomial

$$: u_N^h(x) : = H_h(u_N(x); \sigma_N)$$

ex: Under \mathcal{P} , we have

$$\int_{\mathbb{T}^2} u^2(x) dx = \infty \quad \text{a.s.}$$

$$\text{but } \int_{\mathbb{T}^2} : u_N^2 : = \int_{\mathbb{T}^2} u_N^2 - \sigma_N$$

$$= \sum_{|n| \leq N} \frac{|g_n|^2 - 1}{\langle n \rangle^2}$$

$$\begin{aligned} \text{and } \mathbb{E} \left[\left(\int_{\mathbb{T}^2} : u_N^2 : \right)^2 \right] &= \mathbb{E} \left[(g_0^2 - 1)^2 \right] \\ &\quad + 2 \mathbb{E} \sum_{\substack{n \in \mathbb{Z}^* \\ |n| \leq N}} \frac{(|g_n|^2 - 1)^2}{\langle n \rangle^4} \end{aligned}$$

$$\lesssim 1 + \sum_{|n| \leq N} \frac{1}{\langle n \rangle^4} \lesssim 1 \quad \text{indep of } N$$

$\Rightarrow \int_{\mathbb{T}^2} : u_N^2 :$ is a well-defined r.v.

and we can set

$$\int_{\mathbb{T}^2} : u^2 : = \lim_{N \rightarrow \infty} \int_{\mathbb{T}^2} : u_N^2 :$$

Let

$$G_N(u) = \frac{1}{2m} \int_{\mathbb{T}^2} : (P_{\leq N} u)^{2m} : dx.$$

Prop 2.2: $\{G_N(u)\}$ is a Cauchy seq in $L^q(p)$

for any $q \geq 1$. Moreover, we have

$$\|G_M(u) - G_N(u)\|_{L^q(p)} \leq C_m (q-1)^m \frac{1}{N^{1/2}}.$$

for any $q \geq 1$, $M \geq N \geq 1$.

This proposition allows us to define

$$G(u) = \frac{1}{2m} \int_{\mathbb{T}^2} : u^{2m} : dx$$

$$= \lim_{N \rightarrow \infty} G_N(u) \quad \text{in } L^q(p)$$

and the Wick ordered Hamiltonian

$$H_{\text{Wick}}(u) = \frac{1}{2} \int_{\mathbb{T}^2} |\nabla u|^2 + \frac{1}{2m} \int_{\mathbb{T}^2} : u^{2m} :$$

Pf: (q=2)

$$e_n(y) = e^{in \cdot y}$$

(37)

$$\gamma_N(x)(\cdot) = \frac{1}{\sigma_N^{1/2}} \sum_{|n| \leq N} \frac{\overline{e_n(x)}}{\langle n \rangle} e_n(\cdot)$$

$$\gamma_N(\cdot) = \sum_{|n| \leq N} \frac{1}{\langle n \rangle^2} e_n(\cdot)$$

Note that

$$\|\gamma_N(x)\|_{L^2(\mathbb{T}^2)} = 1 \quad \text{for each fixed } x \in \mathbb{T}^2$$

and

(9) $\langle \gamma_M(x), \gamma_N(y) \rangle_{L^2(\mathbb{T}^2)} = \frac{1}{\sigma_M^{1/2} \sigma_N^{1/2}} \gamma_N(y-x)$

$$\begin{aligned} U_N(x) &= \sigma_N^{1/2} \frac{U_N(x)}{\sigma_N^{1/2}} = \sigma_N^{1/2} \overline{W_{\gamma_N(x)}} \\ &= \sigma_N^{1/2} W_{\gamma_N(x)}. \end{aligned}$$

$$\begin{aligned} \Rightarrow : U_N^{2m}(x) : &= H_{2m}(U_N(x); \sigma_N) \\ &= \sigma_N^m H_{2m}\left(\frac{U_N(x)}{\sigma_N^{1/2}}; 1\right) \\ &= \sigma_N^m H_{2m}(W_{\gamma_N(x)}) \end{aligned}$$

$$\Rightarrow (2m)^2 \|G_M(u) - G_N(u)\|_{L^2(P)}^2$$

$$= \int_{\Pi_x^2 \times \Pi_y^2} \int_{\Omega} \sigma_M^{2m} H_{2m}(W_{\eta_M}(x)) H_{2m}(W_{\eta_M}(y)) \\ - \sigma_M^m \sigma_N^m H_{2m}(W_{\eta_M}(x)) H_{2m}(W_{\eta_N}(y)) \\ - \sigma_M^m \sigma_N^m H_{2m}(W_{\eta_N}(x)) H_{2m}(W_{\eta_M}(y)) \\ + \sigma_N^{2m} H_{2m}(W_{\eta_N}(x)) H_{2m}(W_{\eta_N}(y)) \\ dP dx dy$$

Lemma 2.1

and (9) $= (2m)! \int_{\Pi_x^2 \times \Pi_y^2} (\gamma_M(y-x))^{2m} - (\gamma_N(y-x))^{2m} dx dy$

$$= (2m)! \int_{\Pi^2} \gamma_M(x)^{2m} - \gamma_N(x)^{2m} dx$$

$$\lesssim \int_{\Pi} |\gamma_M(x) - \gamma_N(x)| \left\{ |\gamma_M(x)|^{2m-1} + |\gamma_N(x)|^{2m-1} \right\} dx.$$

$$\bullet \|\gamma_M - \gamma_N\|_{L^2} = \left(\sum_{N < |n| \leq M} \frac{1}{\langle n \rangle^4} \right)^{\frac{1}{2}} \lesssim \frac{1}{N}$$

$$\bullet \|\gamma_N^{2m-1}\|_{L^2} = \|\gamma_N\|_{L^{4m-2}}^{2m-1} \leq \left(\sum_{|n| \leq N} \frac{1}{\langle n \rangle^{\frac{2 \cdot 4m-2}{4m-3}}} \right)^{\frac{4m-3}{2}} \leq C_m < \infty$$

indep of N .



Sec 2.3: Hypercontractivity of the OV semigroup (39)
and Wiener chaos estimate

$$H = L^2(\mathbb{R}^d, \mu_d)$$

↑ standard Gaussian meas on μ_d

$$L = \Delta - x \cdot \nabla$$

= Hartree-Fock operator (Ornstein-Uhlenbeck op)

Lem 2.3 (Hypercontractivity of OV semigroup, Nelson '65)

$p > 1, q \geq 1$. Then, we have

$$\|e^{tL} u\|_{L^q(\mathbb{R}^d, \mu_d)} \leq \|u\|_{L^p(\mathbb{R}^d, \mu_d)}$$

for $t \geq \frac{1}{2} \log\left(\frac{q-1}{p-1}\right)$

Rmk: independent of dimension $d \geq 1$.

Define a homog Wiener chaos of order k

$$H_k(x) = \prod_{j=1}^d H_{k_j}(x_j)$$

$$k = k_1 + \dots + k_d$$

• \mathcal{H}_k = closure of homog Wiener chaoses of order k under $L^2(\mathbb{R}^d, \mu_d)$

\Rightarrow Ito-Wiener decomposition

$$L^2(\mathbb{R}^d, \mu_d) = \bigoplus_{k=0}^{\infty} \mathcal{H}_k$$

$\Rightarrow F \in \mathcal{H}_k$ is an eigenfunction of L with eigenvalue $-k$.

(40)

Cor 2.4: $F \in \mathcal{H}_k$. Then, for $q \geq 2$, we have

$$\|F\|_{L^q(\mathbb{R}^d, \mu_d)} \leq (q-1)^{k/2} \|F\|_{L^2(\mathbb{R}^d, \mu_d)}$$

(\Leftarrow Take $p=2$, $t = \frac{1}{2} \log(q-1)$ in Lemma 2.3.

Lemma 2.5

$$S_k(\omega) = \sum_{\Gamma(k,d)} C(n_1, \dots, n_k) g_{n_1}(\omega) \cdots g_{n_k}(\omega)$$

where $\Gamma(k,d) = \{(n_1, \dots, n_k) \in \mathbb{Z}^k : |n_j| \leq d\}$

\Rightarrow Then, for $q \geq 2$,

$$\|S_k\|_{L^q(\Omega)} \leq \sqrt{k+1} (q-1)^{k/2} \|S_k\|_{L^2(\Omega)}$$

(\Leftarrow Cor 2.4 and $x^{\frac{1}{2}} = \sum_{m=0}^{[\frac{1}{2}]} C_{m,j} \sigma^m H_{j-2m}(x; \sigma)$)

\Rightarrow Prop 2.2 for general $q \geq 2$ follows from the $q=2$ case and Lemma 2.5.

(Note $G_N(u) - G_N(u) \in \bigcup_{k=0}^{2m} \mathcal{H}_k$.)

Sec 2.4: Construction of the Gibbs measure on T^2 .

Construct

$$\begin{aligned} d\mu &= Z^{-1} e^{-H_{\text{wick}}(u)} du \\ &= Z^{-1} e^{-\frac{1}{2m} \int_{\mathbb{T}^2} :u^{2m}: dx} d\rho. \end{aligned}$$

(φ_2^{2m} Euclidean quantum field theory)

$$\text{Let } R_N(u) = e^{-G_N(u)} = e^{-\frac{1}{2m} \int :u_N^{2m}: dx}.$$

Prop 2.6: \bullet $R_N(u) \in L^q(\rho)$ for any $q \geq 1$ with a unif bound in N .

\bullet $R_N(u)$ converges to some $R(u)$ in $L^q(\rho)$.

Pf: $G_N(u)$ is NOT sign definite but.
 - $G_N(u)$ has a logarithmic upper bound:

(10)

$$\begin{aligned} -G_N(u) &= -\frac{1}{2^m} \int_{\mathbb{T}^2} \underbrace{H_{2^m}(u_N; \sigma_N)}_{\geq -a_m} dx \\ &= \sigma_N^m \underbrace{H_{2^m}\left(\frac{u_N}{\sigma_N}\right)}_{\geq -a_m} \\ &\leq b_m (\log N)^m. \end{aligned}$$

Prop 2.2 yields

(11)

$$\begin{aligned} P(q | G_M(u) - G_N(u) | > \lambda) \\ \leq C_m e^{-C_{m,q} N^{\frac{1}{2m}} \lambda^{\frac{1}{m}}} \end{aligned}$$

for all $M \geq N \geq 1$, $q \geq 1$, $\lambda > 0$.

$$\begin{aligned} \bullet \|R_N(u)\|_{L^q(p)}^q &= \int_0^\infty P(e^{-q G_N(u)} > \alpha) d\alpha \\ &\leq 1 + \int_1^\infty P(-q G_N(u) > \log \alpha) d\alpha \end{aligned}$$

Hence, it suffices to show

(12)

$$P(-q G_N(u) > \log \alpha) \leq C \alpha^{-(1+\varepsilon)}$$

for all $\alpha > 1$ and $N \in \mathbb{N}$

Given $\lambda = \log d$,
 choose $N_0 \in \mathbb{R}$ s.t. $\lambda = 2g b m (\log N_0)^m$.

Then, from (10), we have

$$P(-g G_N(u) > \lambda) = 0$$

for all $N < N_0$

• For $N \geq N_0$,

$$P(-g G_N(u) > \lambda)$$

$$\leq P(-g G_N(u) + g G_{N_0}(u) > \lambda - g b m (\log N_0)^m)$$

$$\leq P(-g G_N(u) + g G_{N_0}(u) > \frac{1}{2} \lambda)$$

$$\leq C_m \underbrace{e^{-c} N_0^{1/2m} \lambda^{1/m}}_{e^{-c} \lambda^{1/m} e^{c} \lambda^{1/m}}$$

$$\ll C e^{-(1+\delta)\lambda} \sim \lambda^{-(1+\delta)} \Rightarrow (12)$$

(11) $\Rightarrow G_N(u)$ converges in measure

$\Rightarrow R_N = e^{-G_N(u)}$ converges in measure.

By repeating the argument in Cor 1.4,

$R_N \rightarrow R = e^{-G(u)}$ in $L^q(\mu)$, $\forall q \geq 1$



Rmk: The proof of Prop 2.6 is the so-called Nelson's estimate.

Prop 2.6 allows us to define the Gibbs measure

$$\begin{aligned} d\mu &= Z^{-1} e^{-H_{\text{Wick}}(u)} du \\ &= Z^{-1} e^{-\frac{1}{2m} \int_{\mathbb{R}^2} :u^{2m}:} df. \end{aligned}$$

Sec 2.5: A glimpse of probabilistic local Cauchy theory

• Wick ordered NLS:

$$i\partial_t u + \Delta u = : |u|^{2(m-1)} u :$$

• Wick ordered NLW:

$$-\partial_t^2 u + \Delta u = : u^{2m-1} :$$

• Wick ordered NLKG:

$$-\partial_t^2 u - u + \Delta u = : u^{2m-1} :$$

• Basic idea

• construct (local) solns a.s. with respect to f .

i.e. $u_0 = \sum_{n \in \mathbb{Z}^2} \frac{f_n}{\langle n \rangle} e^{in \cdot x}$

• gain of integrability of lin solns under randomization.

\Rightarrow probabilistic Strichartz estimate

• Write $u = \underbrace{S(t)}_{=Z} u_0 + v$

Z : random, rough (below $L^2(\mathbb{T}^2)$)

v : "deterministic", smoother (in $H^s(\mathbb{T}^2)$ for some $s > 0$)

Bourgain '96: cubic WNLS is a.s. LWP w.r.t. f .

\Rightarrow a.s. GWP & invariance of μ

\uparrow
Case II in Chap 1.

• Also see Colliander - Oh '12.

Rmk: $S_{crit} = 1 - \frac{1}{m-1} \geq \frac{1}{2}$ for (super)quintic $m \geq 3$

\Leftarrow gap is too big.

NLW/NLKG: gain of one derivative in the Duhamel operator. (46)

$$\begin{aligned} : u_N^{2m-1} : &= H_{2m-1}(Z_N + V_N; \sigma_N) \\ &= \sum_{\ell=0}^{2m-1} \binom{2m-1}{\ell} V_N^{2m-1-\ell} H_{\ell}(Z_N; \sigma_N) \end{aligned}$$

$$\text{Scrit} = \max \left(\underbrace{1 - \frac{1}{m-1}}_{\substack{\uparrow \\ \text{from scaling}}}, \underbrace{\frac{3}{4} - \frac{1}{2(m-1)}}_{\substack{\nwarrow \\ \text{from Lorentz inv.}}} \right)$$

Prop 2.7: Probabilistic Strichartz estimates on Wiener homogeneous chaos.

$k \in \mathbb{N}$. Given $2 \leq q, r < \infty$ and $\alpha > 0$,

$$\begin{aligned} &P \left(\| \langle \nabla \rangle^{-\alpha} H_k(Z_N; \sigma_N) \|_{L_T^q L_x^r} > \lambda \right) \\ &\leq C \exp \left(-c \frac{\lambda^2}{T^{1/q}} \right) \end{aligned}$$

for all $T, \lambda > 0$ and $N \in \mathbb{N}$.

Rmk: $k=1 \Rightarrow$ "standard" prob Str. estimate.

$\alpha > 0$ is necessary since $Z_N \in H^{-\varepsilon} \setminus L^2$ a.s.

Idea of Pf: Use the white noise functional ⁽⁴⁷⁾
and the orthogonality relation (Lemma 2.1)

⇒ Theorem 2.8: (Oh-Thomann '15)

WNLW / WNLKG is a.s. LWP w.r.t. ρ .
(\Rightarrow a.s. GWP and invariance of ρ)

Rmk: Due to the lack of mass conservation, we need to consider WNLKG on \mathbb{T}^2 . (or on a domain $M \subset \mathbb{R}^2$ with Neumann B.C.)

On $M \subset \mathbb{R}^2$ with Dirichlet B.C., we can treat WNLW as well.

Sec 2.6: Compactness argument. (Burq-Thomann-Tzvetkov '15)

(Super)quintic WNLS on \mathbb{T}^2 .

$s < 0$: support of the Gibbs meas μ .

$s_{crit} \geq \frac{1}{2}$: scaling critical regularity

⇐ The gap is too big!!

We do not know how to construct strong solutions, even probabilistically.

$$(FWNLS) \quad i\partial_t u^N + \Delta u^N = F_N(u^N)$$

$$\text{where } F^N(u) = P_{\leq N} \left(|P_{\leq N} u|^{2(m-1)} P_{\leq N} u \right)$$

$$= (-1)^{m+1} (m-1)! \sigma_N^{m-1} \cdot P_{\leq N} \left\{ L_{m-1}^{(1)} \left(\frac{|u_N|^2}{\sigma_N} \right) u_N \right\}$$

↑
generalized Laguerre poly.

- μ_N = invariant Gibbs meas for (FWNLS).

① Extend μ_N = prob. meas on initial data

to ν_N = prob meas on space-time functions

- (FWNLS) is globally well-posed.

$$\Rightarrow \exists \Phi_N: u_0 \in H^s \xrightarrow{s \leq 0} u^N \in C(\mathbb{R}; H^s)$$

- Define a prob measure ν_N on $C(\mathbb{R}; H^s)$

$$\text{by } \nu_N = \mu_N \circ \Phi_N^{-1}$$

= induced probability meas under Φ_N

- By invariance of μ_N ,

the law of $u^N(t) = \mu_N$ for any $t \in \mathbb{R}$.

② Show $\{\nu_N\}$ is tight (= compact)

$$\Rightarrow \nu_{N_j} \rightarrow \nu$$

Prokhorov

③ Skorokhod's Thm:

\exists another prob space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$

$\tilde{u}^{N_j}, C(\mathbb{R}, H^s)$ -valued r.v.'s

$u, \quad =$

$$\text{s.t. } \mathcal{L}(\tilde{u}^{N_j}) = \mathcal{L}(u^{N_j}) = \nu_N$$

$$\mathcal{L}(u) = \nu$$

and \tilde{u}^{N_j} converges to u in $C(\mathbb{R}; H^s)$
almost surely w.r.t. \tilde{P} .

Rmk:

$$\mathcal{L}(\tilde{u}^{N_j}(t)) = \mathcal{L}(u^{N_j}(t)) = \mu_N$$

$$\mathcal{L}(u(t)) = \mu \quad (\text{since } \mu_N \xrightarrow{\text{strongly}} \mu)$$

Theorem 2.9 (Oh-Thomann '15)

(50)

\exists a set Σ of full probability s.t.

$\forall \phi \in \Sigma, \exists$ soln $u \in C(\mathbb{R}; H^s)$ to (WNLS)
with $u|_{t=0} = \phi$.

Moreover, $\mathcal{L}(u(t)) = \mu, \forall t \in \mathbb{R}$.

Rmk: only a.s. global existence (NO uniqueness)

$\mathcal{D}_{t,x}$ -valued r.v.

Idea: Let $X_{N_j} = i \partial_t \tilde{u}^{N_j} + \Delta \tilde{u}^{N_j} + F_{N_j}(\tilde{u}^{N_j})$.

Then, we have $\mathcal{L}(X_{N_j}) = \delta_0$

$\Rightarrow \mathcal{L}(X_\infty) = \delta_0$, i.e.

\uparrow
 $u = \lim_{j \rightarrow \infty} \tilde{u}^{N_j}$ is a distr soln to (WNLS)

Need to show

$$: |u|^{2(m-1)} u : = \lim_{N \rightarrow \infty} F_N(u^N)$$

exists in $L^q(\rho; H^s(\mathbb{T}^2))$, $s < 0$.

(analogous to Prop 2.2.

Sec 2.7: Further topics

(51)

① Π^3 :

- Gibbs measure on Π^3 is renormalizable only for $p=4$

- Wick ordering is not enough.

$\sigma_N \sim N$ in 3-d but
need to replace it by $C_1 N + C_2 \log N$.

- stochastic quantization equation:

$$\partial_t u = \Delta u - u^3 + \alpha \cdot u + \xi$$

"local well-posedness".

Hairer '14 (regularity structure)

Catellier-Chouk '14 (paracontrolled distribution)

Rupiainen '14 (renormalization group method)

- NLS/NLKG?

② On \mathbb{R} .

Take the periods $L \rightarrow \infty$ and construct the Gibbs meas on \mathbb{R} as a diffusion process

(Feynmann-Kac formula, Girsanov thm, etc.)

Oh-Quastel-Sosoe '13.

Chap 3: Quasi-invariant Gaussian meas

(52)

$$d\mu_s = Z^{-1} e^{-\frac{1}{2} \|u\|_{H^s}^2} du$$

• μ_s is supported on $H^{s-\frac{d}{2}-\varepsilon}(\mathbb{T}^d)$

$$\omega \mapsto u = \sum_{n \in \mathbb{Z}^d} \frac{g_n}{\langle n \rangle^s} e^{in \cdot x}$$

Q: How is the Gaussian meas μ_s transported under the nonlinear dynamics $\Phi(t): u_0 \mapsto u(t)$?

Is μ_s quasi-invariant?

i.e. μ_s and the pushforward meas $\Phi(t)_* \mu_s = \mu_s \circ \Phi(t)^{-1}$ are equivalent (= mutually abs. conti).

Or, can we prove that μ_s and $\Phi(t)_* \mu_s$ are mutually singular?

Quasi-invariance

• Cameron-Martin '44:

$$\Phi(t) u_0^\omega + \left(\frac{d}{2} + \varepsilon\right) \text{ smoother part indep of } u_0$$

• Ramez '74

$$\Phi(t) u_0^\omega + (d + \varepsilon) \text{ smoother part which may depend on } u_0^\omega$$

Crucial to establish a nonlinear smoothing ⁽⁵³⁾

• Fourth order cubic NLS (cubic biharmonic NLS)
(4NLS) $i\partial_t u = \partial_x^4 u \pm |u|^2 u$ on \mathbb{T}

There is no apparent nonlin. smoothing
 \Rightarrow We use normal form reduction.

Theorem 3.1 : Oh - Tzvetkov '15

Let $s > \frac{3}{4}$. Then, f_s is quasi-invariant
under (4NLS).

Rmk: (4NLS) is globally well-posed in $L^2(\mathbb{T})$
(and ill-posed below L^2), so there is
a gap $\frac{1}{2} < s < \frac{3}{4}$.

• The proof is based on a combination
of global and local analysis.

See the earlier work (Tzvetkov '15) on
the (generalized) BBM equation.

General strategy:

(54)

STEP 1: $\tilde{u}(t) = G[u](t) := e^{2itf|u|^2} u(t)$

$$\Rightarrow i\partial_t \tilde{u} = \Delta_x^4 \tilde{u} + (|\tilde{u}|^2 - 2f|u|^2) \tilde{u}$$

• $v(t) = S(-t) u(t)$ interaction representation.

(13) $\Rightarrow \partial_t v_n = -i \sum_{\Gamma(m)} e^{-i\Phi(\vec{m})} v_n \overline{v_{n_2}} v_{n_3} + i|v_n|^2 v_n$

$$\Gamma(m) = \{ n = n_1 - n_2 + n_3, n_1, n_3 \neq n \}$$

$$\begin{aligned} \Phi(\vec{m}) &= n_1^4 - n_2^4 + n_3^4 - n^4 \\ &= (n_1 - n_2)(n_1 + n_2)(n_1^2 + n_2^2 + n_3^2 + n^2 + 2(n_1 + n_3)^2) \end{aligned}$$

rapid oscillation.

① Show ρ_S is (quasi-) invariant under G and $S(-t)$. Also, under a composition of $T_1 \circ T_2$ where ρ_S is quasi-inv under each T_j .

global
probabilistic

\Rightarrow Then, it suffices to show that ρ_S is quasi-invariant under (13).

STEP 2: Energy estimate.local
PDE analysis

$$\frac{d}{dt} \|v(t)\|_{H^s}^2 = -2 \operatorname{Re} i \sum_n \sum_{l(m)} e^{-i\phi(\bar{n})t} \langle n \rangle^{2s} \overline{v_n} \overline{v_{n_2}} v_{n_3} \overline{v_{n_4}}$$

$$\stackrel{NF}{=} 2 \operatorname{Re} \frac{d}{dt} \left[\sum_n \sum_{l(m)} \frac{e^{-i\phi(\bar{n})t}}{\phi(\bar{n})} \langle n \rangle^{2s} \overline{v_n} \overline{v_{n_2}} v_{n_3} \overline{v_{n_4}} \right]$$

- R :=

$$-2 \operatorname{Re} \sum_n \sum_{l(m)} \frac{e^{-i\phi(\bar{n})t}}{\phi(\bar{n})} \langle n \rangle^{2s} \underbrace{\partial_t (\overline{v_n} \overline{v_{n_2}} v_{n_3} \overline{v_{n_4}})}_{b\text{-lin}}$$

⇒ Defined a modified energy (as in the I-method)

$$E(v) = \|v\|_{H^s}^2 + R(v)$$

Prop 3.2: Let $s > \frac{3}{4}$. $\exists \theta > 0$ s.t.

$$\left| \frac{d}{dt} E(P_{\leq N} v) \right| \leq C \|v\|_{L^2}^{4+\theta} \|v\|_{H^{s-\frac{1}{2}-\varepsilon}}^{2-\theta}$$

STEP 3: Construction of weighted Gaussian meas

$$d\mu_{s,N,r} = Z_{s,N,r}^{-1} \mathbb{1}_{\{\|v\|_{L^2} \leq r\}} e^{-\frac{1}{2} R(P_{\leq N} v)} df_s$$

$$d\mu_{s,r} = Z_{s,r}^{-1} \mathbb{1}_{\{\|v\|_{L^2} \leq r\}} e^{-\frac{1}{2} R(v)} df_s$$

• analogous to Chap 1.

STEP 4: change-of-variable formula

(56)

$$d\mu_{s,N,r} = Z_{s,N,r}^{-1} \mathbb{1}_{\{\|v\|_{L^2} \leq r\}} e^{-\frac{1}{2} E(P_{\leq N} v)} dL_N \otimes df_{s,N}^\perp$$

global

$$dL_N = \prod_{|n| \leq N} d\hat{u}_n = \text{Leb meas on } \mathbb{C}^{2N+1}$$

Prop 3.3 $s > 1/2$, $N \in \mathbb{N}$, $r > 0$. Then,

$$\mu_{s,N,r}(\Psi_N(t) A)$$

$$= Z^{-1} \int_{\Psi_N(t) A} \mathbb{1}_{\{\|v\|_{L^2} \leq r\}} e^{-\frac{1}{2} R(P_{\leq N} v)} df_s(v)$$

$$= Z^{-1} \int_A \mathbb{1}_{\{\|v\|_{L^2} \leq r\}} e^{-\frac{1}{2} E(P_{\leq N} \Psi_N(t) v)} dL_N \otimes df_{s,N}^\perp.$$

$\Psi_N(t)$ = soln map to

$$\partial_t v_n = \mathbb{1}_{|n| \leq N} \left\{ -i \sum_{\substack{|m| \leq N \\ |n_j| \leq N}} e^{-i\phi(n)t} v_n v_{n_2} v_{n_3} + i |v_n|^2 v_n \right\}$$

• L_N is invariant under $P_{\leq N} \Psi_N(t) P_{\leq N}$

low freq dynamics

STEP 5: evolution of truncated measures

(57)

Lemma 3.4: let $s > \frac{3}{4}$. Then, $\exists \beta \in [0, 1)$

s.t. $\forall r \exists C > 0$ s.t.

$$\left| \frac{d}{dt} \mu_{s, N, r}(\Psi_N(t)(A)) \right| \leq C \left\{ \mu_{s, N, r}(\Psi_N(t)(A)) \right\}^{1-\frac{1}{p}}$$

for any $p \geq 2$, $N \in \mathbb{N}$, meas set $A \subset L^2(\mathbb{T})$

Pf: Idea: Reduce the problem to $t=0$

$$\left| \frac{d}{dt} \mu_{s, N, r}(\Psi_N(t)(A)) \right|_{t=t_0}$$

$$= \left| \frac{d}{dt} \mu_{s, N, r}(\Psi_N(t)(\Psi_N(t_0)(A))) \right|_{t=0}$$

Prop 3.3

$$= \mathbb{E} \left[\frac{d}{dt} \int_{\Psi_N(t_0)(A)} \mathbb{1}_{\{\|w\|_{L^2} \leq r\}} e^{-\frac{1}{2} E(\underline{P}_N \Psi_N(t)(w))} dL_N \otimes df_{s, N}^1 \right]_{t=0}$$

$$= -\frac{1}{2} \int_{\Psi_N(t_0)(A)} \left| \frac{d}{dt} E(\underline{P}_N \Psi_N(t)(w)) \right|_{t=0} d\mu_{s, N, r}$$

Now, apply Energy estimate (Prop 3.2)
with Hölder's ineq.

□

• Lemma 3.4 and Yudovich's argument

⇒ Lemma 3.5: Let $s > 3/4$. Given $t \in \mathbb{R}$, $r > 0$, and $\delta > 0$, $\exists C = C(t, r, \delta) > 0$ s.t.

$$\mu_{s, N, r}(\Psi_N(t)(A)) \leq C \{\mu_{s, N, r}(A)\}^{1-\delta}$$

for any $N \in \mathbb{N}$ and measurable $A \subset L^2(\mathbb{T})$

STEP 6: Quasi-invariance of ρ_s .

• Lemma 3.5 and approximation argument

⇒ Lemma 3.6: Let $s > \frac{3}{4}$. Given $t \in \mathbb{R}$, $r > 0$, and $\delta > 0$, there exists $C = C(t, r, \delta) > 0$ s.t.

$$\mu_{s, r}(\Psi(t)(A)) \leq C \{\mu_{s, r}(A)\}^{1-\delta}$$

↑
soln map for (13). on page (54)

Pf of Thm 3.1: Suppose $\rho_s(A) = 0$.

⇒ Then, $\rho_{s, r}(A) = 0$, where $d\rho_{s, r} = \mathbb{1}_{\{\|u\|_{L^2} \leq r\}} d\rho_s$ for any $r > 0$.

⇒ $\mu_{s, r}(A) = 0$ b/c $\mu_{s, r} \ll \rho_{s, r}$

⇒ By Lemma 3.6, $\mu_{s, r}(\Psi(t)(A)) = 0$.

⇒ $\rho_{s, r}(\Psi(t)(A)) = 0$ b/c $\rho_{s, r} \ll \mu_{s, r}$.

$$\Rightarrow p_s(\Psi(t)A) \stackrel{\text{DCT}}{=} \lim_{r \rightarrow \infty} p_{s,r}(\Psi(t)A) = 0.$$

□

Rmk: This argument allows us to obtain

$$\|u(t)\|_{H^\sigma} \lesssim t^{\alpha(\sigma)} \quad \text{for } \sigma < s - \frac{1}{2}$$

by adapting Bourgain's argument on p. (24)